

# Know your footprint - in High Energy Physics and related fields


Valerie Lang

On behalf of the Know-your-footprint team

Naman Kumar Bhalla, Simran Gurdasani, Pardis Niknejadi, VL

Particle Physics Colloquium

Paul Scherrer Institute, 24 October 2024



# Atmospheric CO<sub>2</sub> vs. ground temperature

First publication on their relationship in 1896, i.e. 128 years ago

- Prof. Svante Arrhenius, *On the Influence of Carbonic Acid in the Air upon the Temperature of the Ground* Philosophical Magazine and Journal of Science Series 5, Volume 41, April **1896**, pages 237-276. ([link](#))

TABLE VII.—Variation of Temperature caused by a given Variation of Carbonic Acid.

Europe

Latitude.	Carbonic Acid=0.67.						Carbonic Acid=1.5.						Carbonic Acid=2.0.						Carbonic Acid=2.5.						Carbonic Acid=3.0.					
	Dec.-Feb.	March-May.	June-Aug.	Sept.-Nov.	Mean of the year.		Dec.-Feb.	March-May.	June-Aug.	Sept.-Nov.	Mean of the year.		Dec.-Feb.	March-May.	June-Aug.	Sept.-Nov.	Mean of the year.		Dec.-Feb.	March-May.	June-Aug.	Sept.-Nov.	Mean of the year.		Dec.-Feb.	March-May.	June-Aug.	Sept.-Nov.	Mean of the year.	
70	-2.0	-3.0	-3.4	-3.1	-3.1		3.3	3.4	3.8	3.6	3.3		6.0	6.1	6.0	6.1	6.05		7.9	8.0	7.9	8.0	7.95		9.1	9.3	9.4	9.4	9.3	
60	-3.0	-3.2	-3.4	-3.3	-3.22		3.4	3.7	3.6	3.8	3.62		6.1	6.1	5.8	6.1	6.02		8.0	8.0	7.6	7.9	7.87		9.3	9.5	8.9	9.5	9.3	
50	-3.2	-3.3	-3.3	-3.4	-3.3		3.7	3.8	3.4	3.7	3.65		6.1	6.1	5.5	6.0	5.92		8.0	7.9	7.0	7.9	7.7		9.5	9.4	8.6	9.2	9.17	
40	-3.4	-3.4	-3.2	-3.3	-3.32		3.7	3.8	3.3	3.6	3.55		6.1	6.1	5.1	5.6	5.71		8.0	7.9	6.8	7.6	7.42		9.5	9.4	8.2	9.2	9.02	
30	-3.3	-3.2	-3.1	-3.1	-3.17		3.5	3.3	3.2	3.5	3.37		5.6	5.4	5.0	5.2	5.3		7.2	7.0	6.6	6.7	6.87		8.7	8.3	7.5	7.9	8.1	
20	-3.1	-3.1	-3.0	-3.1	-3.07		3.5	3.2	3.1	3.2	3.25		5.2	5.0	4.9	5.0	5.02		6.7	6.6	6.3	6.6	6.52		7.9	7.5	7.2	7.5	7.52	
10	-3.1	-3.0	-3.0	-3.0	-3.02		3.2	3.2	3.1	3.1	3.13		5.0	5.0	4.9	4.9	4.95		6.6	6.4	6.3	6.4	6.42		7.4	7.3	7.2	7.3	7.3	
0	-3.0	-3.0	-3.1	-3.0	-3.02		3.1	3.1	3.2	3.2	3.13		4.9	4.9	5.0	5.0	4.95		6.4	6.4	6.6	6.6	6.5		7.3	7.3	7.4	7.4	7.35	
-10	-3.1	-3.1	-3.2	-3.1	-3.12		3.2	3.2	3.2	3.2	3.17		5.0	5.0	5.2	5.1	5.07		6.6	6.6	6.7	6.7	6.63		7.4	7.5	8.0	7.6	7.62	
-20	-3.1	-3.2	-3.3	-3.2	-3.2		3.2	3.2	3.4	3.3	3.27		5.2	5.3	5.5	5.4	5.35		6.7	6.8	7.0	7.0	6.87		7.9	8.1	8.6	8.3	8.22	
-30	-3.3	-3.3	-3.4	-3.4	-3.35		3.4	3.5	3.7	3.5	3.52		5.5	5.6	5.8	5.6	5.62		7.0	7.2	7.7	7.4	7.32		8.6	8.7	9.1	8.8	8.8	
-40	-3.4	-3.4	-3.3	-3.4	-3.37		3.6	3.7	3.8	3.7	3.65		5.8	6.0	6.0	6.0	5.95		7.7	7.9	7.9	7.9	7.85		9.1	9.2	9.4	9.3	9.25	
-50	-3.2	-3.3	—	—	—		3.8	3.7	—	—	—		6.0	6.1	—	—	—		7.9	8.0	—	—	—		9.4	9.5	—	—	—	
-60	—	—	—	—	—		—	—	—	—	—		—	—	—	—	—		—	—	—	—	—		—	—	—	—	—	




- CO<sub>2</sub> increase by a factor 2: Temperature increase of ~6°C
- Surprisingly accurate given coarse understanding at the time

- Confirmed and refined since then in many studies → See e.g. Nobel prize in 2021

# Guess the concentration of CO<sub>2</sub> in the atmosphere

What is the current CO<sub>2</sub> content in the atmosphere, given a mean over the last 800k years of around 225 ppm?



Or go to  
[www.vote.ac](http://www.vote.ac)  
and type my  
email address


Reset Vote

ID = valerie.lang@physik.uni-freiburg.de  
36 participants / Poll closed

# Where are we now? – In terms of CO<sub>2</sub> in atmosphere

Measurements over the last ~70 years at Mauna Loa Observatory → Keeling curve

- Combined with data from ice cores over last 800k years → Composition of air trapped in ice from Antarctica



<https://keelingcurve.ucsd.edu>




# Where are we now? – In terms of ground temperature

Copernicus Satellite Data → Sept. 2024 = 2<sup>nd</sup> warmest on record (after Sept. 2023)

Surface air temperature anomaly for September 2024

Data: ERA5. Reference period: 1991-2020. Credit: C3S/ECMWF



PROGRAMME OF  
THE EUROPEAN UNION




IMPLEMENTED BY



Daily sea surface temperature for 60°S–60°N

Data: ERA5 1979–2024 • Credit: C3S/ECMWF



PROGRAMME OF  
THE EUROPEAN UNION



- Over the last 12 months (from September 2024):
  - Average global temperature: 1.62°C above 1850-1900 level
  - Average European temperature: 2.3°C above 1850-1900 level

<https://climate.copernicus.eu/surface-air-temperature-september-2024>

<https://climate.copernicus.eu/climate-bulletin-about-data-and-analysis>

Reached (and surpassed) the global 1.5°C increase limit, ideally targeted by Paris climate agreement 2015

# Intergovernmental Panel on Climate Change (IPCC)

Comprehensive reports on the state of climate change, its impacts and risks, as well as mitigation strategies → Latest: Sixth Assessment report (AR6)

- Working Group I – The Physical Science Basis → Released Aug 2021
- Working Group II – Impacts, Adaptation and Vulnerability → Released Feb 2022
- Working Group III – Mitigation of Climate Change → Released April 2022
- Synthesis Report → Released March 2023



Nobel

2007 PEACE PRIZE  
© THE NOBEL FOUNDATION

→ From the Summary for Policy Makers of the Synthesis Report:

**A.1 Human activities, principally through emissions of greenhouse gases, have unequivocally caused global warming, with global surface temperature reaching 1.1°C above 1850-1900 in 2011-2020. Global greenhouse gas emissions have continued to increase, with unequal historical and ongoing contributions arising from unsustainable energy use, land use and land-use change, lifestyles and patterns of consumption and production across regions, between and within countries, and among individuals (*high confidence*). {2.1, Figure 2.1, Figure 2.2}**

<https://www.ipcc.ch/reports/>

[https://www.ipcc.ch/report/ar6/syr/downloads/report/IPCC\\_AR6\\_SYR\\_SPM.pdf](https://www.ipcc.ch/report/ar6/syr/downloads/report/IPCC_AR6_SYR_SPM.pdf)

# Where are we heading?


Also see IPCC WGI

Interactive Atlas:


<https://interactive-atlas.ipcc.ch>

## Different scenarios in IPCC report analysed

- Factoring (lack of) mitigation actions, policies, etc.



- Pathways to 1.5°C (2.0°C) require rapid and deep yearly emissions reductions!
- Why? Cumulative CO<sub>2</sub> emissions count



→ Currently implemented policies lead to warming of 3.2°C

[https://www.ipcc.ch/report/ar6/syr/downloads/report/IPCC\\_AR6\\_SYR\\_SPM.pdf](https://www.ipcc.ch/report/ar6/syr/downloads/report/IPCC_AR6_SYR_SPM.pdf)


[https://report.ipcc.ch/ar6syr/pdf/IPCC\\_AR6\\_SYR\\_LongerReport.pdf](https://report.ipcc.ch/ar6syr/pdf/IPCC_AR6_SYR_LongerReport.pdf)



# Why is $> 2.0^{\circ}\text{C}$ temperature increase a bad idea?

With warming of  $3.2^{\circ}\text{C}$ :

- ~100% biodiversity losses in large areas near equator
- Large parts of the Earth become ~uninhabitable due to risk of hyperthermia



→ Hyperthermia = Failure of human heat-regulating mechanisms – deadly if not treated quickly




# Why is it relevant to High Energy Physics & related fields?

---

## High Energy Physics (HEP) and related fields contribute to CO<sub>2</sub> emissions

- Build large detector systems and infrastructures
  - Cause emissions from various sources  
→ See environmental reports e.g. by CERN
- But: How much per researcher? → **Know your footprint!**
  - Idea: Estimate per-researcher carbon footprint  
→ Put into context with private and target footprints
  - Personal identification of high-emission areas which need urgent addressing and raise awareness
  - Provide personal reference for gauging carbon emission numbers



→ If we want to maintain ~liveable conditions on Earth, ALL areas of research, politics, culture, industry, etc. need to contribute to emissions reductions → This includes HEP!

# Know your footprint (Kyf) calculator

## Consider private and professional emissions for researchers

- Private emissions in Germany – see [carbon calculator](#) by German Federal Environment Agency (UBA)
- Professional emissions in HEP and related fields
  - Split into four categories:
    - Experiment
    - Institute
    - Computing
    - Travel

→ Investigate each category's impact  
→ Configurable per individual researcher, i.e. your individual research situation!

→ Know your footprint (Kyf) calculator  
<https://limesurvey.web.cern.ch/863499?lang=en>


→ Paper discussing the basis of the Kyf calculator  
<https://arxiv.org/abs/2403.03308>

→ Discuss in the following

### Carbon footprints in comparison

Carbon footprint: 10.13 t

German average: 10.35 t



# Experiment, collaboration or project footprint

## Distinguish the following options

- Large LHC experiment
  - Small LHC experiment
  - Small HEP experiment
  - Astrophysics experiment
- Based on CERN environmental report(s)
- Based on DESY electricity consumption
- Based on ESO annual report → Skip today



## Definition of per-researcher footprint per year


- Per-researcher footprint = (Total annual emissions from experiment) / (Number of experiment members)
  - Experiment members = collaboration members or users (and operators) according to applicability
  - No consideration of indirect benefits for “the industry” or “the public” through “gained knowledge”  
→ Too vague and leads to responsibility diffusion
  - Responsibility for emissions lies with researchers designing, building, and operating detectors, and analyzing their data

# Footprint of large and small LHC experiments

## Emissions classified into three categories by CERN environmental reports

- Scope 1
  - Direct emissions from detectors, heating, etc.
- Scope 2
  - Indirect emissions, primarily from electricity consumption
- Scope 3 → Considered only for Institute footprint
  - Indirect emissions from other sources, e.g. travel, commute, waste, catering, procurement

→ Average emissions separately over:  
Running years: 2017, 2018, 2022, and  
Shutdown years: 2019-2021





# Footprint of large and small LHC experiments (II)

## Assign emissions to large and small LHC experiments

- For scope 1 emissions → LHCb Scope 1 emissions in 2022 specified in Upgrade II Technical Design report
  - Assume ALICE ≈ LHCb → Small LHC experiment:  $S1_{small}$
  - Assume ATLAS ≈ CMS → Large LHC experiment:  $S1_{Large}$


$$\rightarrow S1_{Large} = \frac{S1_{All} - 2 \cdot S1_{small}}{2}$$

- For scope 2 emissions
  - Largest consumer: LHC → Followed by pre-accelerators  
→ Needed by all four experiments → Share equally  
→ Subtract CERN-site contributions before

$$\rightarrow S2_{Large} = S2_{small} = \frac{S2_{All}}{4}$$

- Typical operation pattern in last years: 4 years of running, 3 years of shutdown
  - Weight accordingly for overall annual emissions → Total emissions per experiment

Electrical power distribution 2018



# Footprint of large and small LHC experiments (III)

Calculate per-researcher emissions per year

- Assume equal share among collaboration members

Emissions [tCO<sub>2</sub>e] per experiment

	Phase	Scope 1	Scope 2	Total
Small	Run	2244	16 206	18 450
	SD	1030	8796	9826
	Overall	-	-	14 754
Large	Run	78 332	16 206	94 538
	SD	35 962	8796	44 758
	Overall	-	-	73 204

Emissions per collaboration member

	Experiment	Members	Mean	Emissions
Small	ALICE	1968	1684	8.76 tCO <sub>2</sub> e
	LHCb	1400		
Large	CMS	6288	6144	11.91 tCO <sub>2</sub> e
	ATLAS	6000		


→ Slightly more (less) for large (small) LHC experiments compared to the private footprint in Germany

# Footprint of a small HEP experiment

## Estimate based on DESY electricity consumption

- Data from 2021: 128.3GWh annually
- Convert to tCO<sub>2</sub>e → 2 options:
  - Green electricity  
→ Assume 100% photo-voltaic (PV) based production → 35 gCO<sub>2</sub>e/kWh
  - German electricity mix in 2023  
→ Includes >40% from wind, solar and water power → 416 gCO<sub>2</sub>e/kWh (for comparison: gas: 572 gCO<sub>2</sub>e/kWh, coal: 1167 gCO<sub>2</sub>e/kWh)

### Energy consumption DESY 2021



→ With 3000 guest scientists + 200 operators: 1.40 tCO<sub>2</sub>e (16.68 tCO<sub>2</sub>e) with green (conventional) electricity

# Institute or research centre footprint

---

## Distinguish the options

- University (with green or conventional electricity)
  - Based on University of Freiburg report (skip Leibniz University Hannover today)
- Research centre
  - Based on CERN environmental report(s)



## Definition of per-researcher footprint per year


- $(\text{Total institute emissions}) / (\text{Effective number of institute members})$ 
  - One representative year outside of COVID-19 pandemic: 2019 for University of Freiburg, 2022 for CERN
- University of Freiburg as default university footprint
  - Omission of procurement information by Leibniz University Hannover
  - Decent agreement in overlapping categories between both universities



# Footprint of a university - Freiburg


## Emissions with green electricity

- Exclude emissions from travel here



## Procurement → Dominating contributor

- Based on procurement data from 2017




→ Many categories → Challenging to address  
→ Demand management + green procurement!

# Footprint of a university - Freiburg

## Emissions with green electricity

- Exclude emissions from travel here



Increases to  
19224 tCO<sub>2</sub>e  
with conventional  
electricity


→ With 31147 members of the university (students and employees):  
1.00 tCO<sub>2</sub>e (1.54 tCO<sub>2</sub>e) with green (conventional) electricity

\* Die Emissionen aus der Beschaffung basieren auf den Ausgaben des Jahres 2017

\*\* Die Emissionen durch Dienstreisen basieren auf einer Hochrechnung für das Jahr 2018

## Procurement → Dominating contributor

- Based on procurement data from 2017




→ Many categories → Challenging to address  
→ Demand management + green procurement!

# Footprint of a research centre – CERN


## From the CERN environmental reports

- Heating + Other category from scope 1
- 5% of electricity, i.e. scope 2
- Scope 3
  - Excluding commute, travel, and catering



## Procurement contribution = huge!

- Procurement emissions: 104 974 tCO<sub>2</sub>e in 2022!
- Corresponds to ~57% of total scope 1 emissions in same year
- Contributions for construction of future infrastructure, etc. included → Cannot be clearly separated → Maintain fully under institute



EMISSIONS BY PROCUREMENT FAMILY 2021–2022

"Other" includes: office supplies, furniture, transport, handling and vehicles; centralised expenses and codes for internal use; particle and photon detectors; health, safety and environment; optics and photonics.

# Footprint of a research centre – CERN (II)

## Total institute emissions

Category	Emissions [tCO <sub>2</sub> e ]
Electricity	3158
Heating (gas+fuel) + Other	11 250
Water purification	176
Waste	1875
Procurement	104 974
Total	121 433
Total without Procurement	16 459

## Effective CERN population

- At any time during the year:
  - Fraction of CERN users at CERN, using electricity, heating, water, etc.
  - Consider together with CERN personell, i.e. staff and CERN fellows

→ Effective CERN population: 7295

→ Per-researcher footprint:  
16.65 tCO<sub>2</sub>e (2.26 tCO<sub>2</sub>e) including (excluding) procurement  
→ With procurement, artifically increased, due to impossibility of procurement split-up  
→ Needs update, once more refined data available

→ To CERN's credit:

Environmentally Responsible Procurement Policy, effective from 1 January 2024 – [April 2024 CERN news](#)

→ Hopefully, procurement footprint will reduce over the next years



# Computing footprint

## Focus on High Performance Computing (HPC)

- Specify individual's computing workloads in core hours
- Distinguish between CPU and GPU usage
  - Choice of CPU or GPU due to computational task
    - Several possibilities to tune configuration
    - Assume optimal core utilization
- Possibility to add footprint of large external (commercial) data storage resources
- Personal computers, small institute clusters, etc. not included
  - Assumed to be covered by personal or institute electricity bills and procurement → Thus included in personal or institute footprint
- Four benchmark scenarios for easy use available



# Computing footprint (II)

---

## Calculation of computing footprint

$$Total [tCO_2e] = f_{PUE} \cdot f_{overh} \cdot n_{WPC} \cdot f_{conv}$$

- With:

- $f_{PUE}$  = HPC's Power Usage Effectiveness (PUE)  
→ Default: 1.5 (Global average) → New CERN computing centre target: 1.1 ([Feb 2024 CERN news](#))
- $f_{overh}$  = Overhead factor for power consumption when computing cores are idle  
→ Default: 1.17 (Hawk supercomputer idle time at the HPC Stuttgart)
- $n_{WPC}$  = Workload Power Consumption (WPC)

$$n_{WPC} = p_{CPU-core} \cdot l_{core-h,CPU} + p_{GPU} \cdot l_{h,GPU}$$

$p_{CPU-core/GPU}$  = Power consumption in kW for each CPU core/GPU

→ Default: 7.25W (CPU - from the DESY Maxwell cluster with AMD EPYC 75F3 CPU cores),  
250W (GPU - median of range, reported on a forum of NVIDIA GPU users)

$l_{core-h,CPU/h,GPU}$  = CPU workload measured in core hours/ GPU usage hours → User input

- $f_{conv}$  = Conversion factor from kWh to gCO<sub>2</sub>e → Both, green and conventional (default) electricity possible

# Computing footprint (III)

---

## Four benchmark scenarios

- Low usage
  - PhD student with several jobs per week → Average of 4000 CPU core-h/month
- Medium usage
  - Doctoral student or post-doctoral researcher, strongly involved in data analysis → Based on top five ranked users at the Uni-Freiburg HPC: Black-Forest Grid (BFG) → Average of 30 000 CPU core-h/month
- High usage
  - Accelerator scientist, studying accelerator performance with particle tracking codes and semi particle in-cell (PIC) codes → With code optimized for GPUs: 2500 GPU h/month ( $\approx$  80 000 CPU core-h/month)
- Extremely high usage
  - Researcher running PIC simulations or high-resolution imaging analysis → 8000 GPU h/month ( $\approx$  300 000 CPU core-h/month)

With conventional electricity	
Scenario	Annual footprint [tCO <sub>2</sub> e]
Low	0.25
Medium	1.91
High	5.48
Extremely high	17.52

# Travel

---

Consider only business travel → Private travel included in private footprint

- Travel important in international research environment:
  - For personal connections at in-person meetings
  - For building research networks, collaborations
  - Etc.
- Most notably missed during COVID-19 pandemic
- BUT: Travel creates CO<sub>2</sub> emissions
  - Which travel is essential and which is not?
  - Re-evaluate how travel is performed:
    - Longer travel times with non-air based travel  
= longer-duration stays preferable
    - Constraints from teaching, family, etc. = non-trivial



Foto von [detail](#) auf [Unsplash](#)

- Possibility for detailed calculations of business trip emissions in Kyf calculation OR benchmark trips



# Travel (II)

## Based on information from the German UBA

- German numbers for hotel and venue assumed to be valid internationally

Source of Emission	Emission Factor	
Long-distance Buses	0.031	kgCO <sub>2</sub> e/km
Long-distance Trains	0.031	kgCO <sub>2</sub> e/km
Personal Car	0.17	kgCO <sub>2</sub> e/km
Flights within Europe	130	kgCO <sub>2</sub> e/h
Transcontinental Flights	170	kgCO <sub>2</sub> e/h
Hotel room	12	kgCO <sub>2</sub> e/night
Event venue	0.19	kgCO <sub>2</sub> e/day




Benchmark	Situation	Emissions [tCO <sub>2</sub> e]
Travel within Germany	5-day trip by trains from Freiburg to Hamburg	0.1
	Same but by plane (1.5h flight/direction)	0.5
Travel within Europe	5-day trip by plane from Freiburg to Thessaloniki (2.5h flight/direction)	0.7
Travel across continents	2-week trip by plane from Freiburg to Seoul (12h flight/direction)	4.3

- In particular, cross-continental flights contribute significantly
- CO<sub>2</sub> compensation for flights possible to indicate in Kyf calculator

# Benchmark researcher

## Putting everything together

- Benchmark for early-career researcher in Germany: Doctoral student
    - Working on one of the large LHC experiments
    - Employed by university with conventional electricity
    - Medium computing level with conventional electricity
    - Annual travel: Two 1-week trips by train in Germany, one 1-week flight travel in Europe, 1 2-week cross-continental travel (e.g. for summer school)
- Professional footprint exceeds private footprint by factor of ~2
- Both by far exceed targets for mitigating climate crisis to only 2.0°C or 1.5°C warming
- HEP research urgently needs to address this
- Become part of the solution of the climate crisis!



# Summary

---

## Climate crisis in progress and intensifying every year

- Mechanism of CO<sub>2</sub> concentration and ground temperature increase known since more than 100 years
    - Currently heading towards 3.2°C temperature increase → Will cause in some areas ~100% biodiversity loss, and makes large regions on the planet deadly for human life
    - Targeted action for mitigation urgently needed!
  - High Energy Physics (HEP) and related areas contribute to global emissions → Reductions urgent
    - Know your footprint (Kyf) calculator for individual researcher emissions
    - Evaluation for early-career benchmark researcher: Professional and private footprint together factor of ~6 (~18) larger than needed for 2.0°C (1.5°C) temperature increase mitigation
      - Every gram of CO<sub>2</sub> not emitted counts!
- Know your footprint to know where to start! → If large contributions from:
- Experiment and institute: Send e-mail to your experiment and institute responsables asking for details
  - Computing: Send e-mail to your computing center about using renewables, think about efficient coding
  - Travel: Think about which travel is needed and which means is possible

# Take-away from today

What is your most important take-away from this seminar today? (1-2 words max.)

Word cloud content:

- Will not help if corporation...
- Very usefull, important, ple...
- Act now
- collective effort.
- Doomed
- I didn't know that we can di...
- PSI, start firing
- professional footprint close...
- Missing data
- Next generation
- SAD
- CO2 footprint
- Interesting and surprising r...
- Shocking
- Too much cooling
- More efficient code
- Global efforts
- Reflect.
- Hopeless
- The only solution is to brea...
- Large professional emissions
- impossible to solve
- Detector Cooling
- Don't use Deutsche Bahn
- What about PSI??
- Nuclear power plants are the...
- Hypothermia scary
- Shut down CERN to stop CO2 é...
- Track not only personal but ...
- 2 deg not possible any more?



Reset Vote

ID = [valerie.lang@physik.uni-freiburg.de](mailto:valerie.lang@physik.uni-freiburg.de)  
36 Posts / Poll closed



# Know your footprint! – Questions?

Thanks for your attention

→ If possible, please submit your data (anonymously) so that we can get an overview of the averages



# Sustainability at PSI

---

## PSI environmental mission statement and PSI Energy mission statement

- Environmental performance based on the following indicators
    - Efficient energy use in infrastructure and research (see PSI Energy Mission Statement)
    - Reduction of the emissions caused directly and indirectly
    - Optimised use of resources (water, paper, chemicals etc.)
    - Advanced waste management with a high proportion of recycling
  - Also annual reports
    - PSI submits a comprehensive and transparent annual report to the ETH Board on key figures, measures and the achievement of objectives
  - Core objectives
    - Increasing energy efficiency
    - Expansion of green power generation
    - Reduction of direct and indirect emissions
      - e.g. reduction of 50% of greenhouse gas emissions from 2006 to 2030
    - Efficient use of resources (water, paper, chemicals, etc.)
    - Progressive waste management with a high proportion of material recycling
- Two very important steps for emission reductions (already):
- Since 2020, PSI has been obtaining all its electricity with hydropower certificates and implements the targets for increased electricity generation with photovoltaics.
  - PSI specifically advocates the operation of the new comprehensive heat recovery system and its expansion.
- More numbers in ETH and VKE reports (see next slides)

# Sustainability at PSI (II)

## In ETH Domain report 2023

- Page 108
- Extracted PSI information only

→ Due to temporarily increased energy prices for hydro power (electricity shortage), nuclear power had to be used temporarily, listed under uncertified electricity

PSI		
<b>BASIC DATA</b>		
Energy reference area (ERA) <sup>2</sup>	m <sup>2</sup>	172,571
Full-time equivalent <sup>3</sup>	FTE	2,132
<b>ENERGY<sup>4</sup></b>		
Final energy, net <sup>5</sup>	kWh/a	144,800,906
<b>Electricity, net (not incl. self-produced)</b>		
Consumption of uncertified electricity	kWh/a	3,024,646
Consumption of certified electricity	kWh/a	135,296,354
Electricity (without naturemade star)	kWh/a	135,000,000
Photovoltaic naturemade star	kWh/a	296,354
Hydro power naturemade star	kWh/a	0
Wind naturemade star	kWh/a	0
Sale of electricity	kWh/a	0
<b>Heat</b>		
Fuel oil	kWh/a	5,390,430
Natural gas, biogas	kWh/a	240,430
District heating	kWh/a	0
Woodchip	kWh/a	5,150,000
Sale of heat	kWh/a	0
<b>Fuel (own vehicles)</b>	kWh/a	189,476
<b>Energy: additional information</b>		
Energy costs, electricity and heat <sup>6</sup>	CHF/a	15,786,345
Self-generated renewable electricity	kWh/a	354,028
Total sale to third parties	kWh/a	0
<b>WATER (DRINKING WATER)</b>		
	m <sup>3</sup>	63,865
<b>MATERIALS</b>		
Paper	kg	18,106
Paper, new fibre	kg	10,322
Paper, recycled	kg	7,784
<b>KEY FIGURES: ENVIRONMENTAL IMPACT</b>		
Primary energy <sup>4</sup>	kWh/a	181,473,226
Proportion of renewable energies	%	89
CO <sub>2</sub> emissions <sup>8</sup>	t CO <sub>2</sub> /a	595

- Provisional figures for the year under review (trend), as of: start of March 2024.
- The energy reference area is the sum of all gross floor areas, above and below ground, that must be heated or air-conditioned in order to be used.
- The FTE (full-time equivalent) value listed here was supplemented by the number of students with an FTE value of 0.68 to produce the consumption per person.
- The key figures indicated for electricity and heat show the total consumption of both for buildings, as well as for teaching and research activities.
- The key indicator "energy costs" shows all expenditure (cash out) for the provision of energy (heat and electricity).
- In energy economics, one refers to primary energy as the energy that is available using the original forms or resources of energy, such as fuel (e.g. coal or natural gas), as well as energy carriers such as sun, wind or nuclear fuels.
- Final energy is the portion of the primary energy that is left after losses due to energy conversion and transmission after it is supplied via the consumer's domestic connection. The final energy basically corresponds to the energy that is purchased.
- CO<sub>2</sub> emission factors according to Ecoinvent version 3.71.

→ Total of 595 tCO<sub>2</sub>e/year → With 2132 FTE (assume here = number of people – not fully true – see above) → **~0.3 tCO<sub>2</sub>e/year/person**  
 → However, only "Paper" seems included for procurement – procurement at CERN or University of Freiburg significant contribution

## Also interesting related to PSI


- Swiss Center for Excellence on Net zero Emissions (SCENE)
- More information available for PSI employees
  - <https://intranet.psi.ch/de/uem>
  - <https://intranet.psi.ch/de/dir/news/neujahrsinformation-fuer-mitarbeitende-vom-16-januar-2024>

# Sustainability at PSI (III)

## Schweizerische Eidgenossenschaft Vorbild Energie und Klima (VEK) → Incl. PSI

- In VEK energy and climate report 2023


- Final energy usage in 2023



→ Includes PSI

### Treibhausgasemissionen

For ETH-area (incl. PSI)



→ Individual institute measures (PSI and all ETH)

**5. CO<sub>2</sub>-Emissionen aus Geschäftsflügen reduzieren**

■ -30 % (2030)

**7. Synchrotron Lichtquelle Schweiz (SLS) 2.0 (PSI)**

■ -2 GWh/a (2025)



# Carbon footprint Switzerland

## Per capita CO<sub>2</sub> emissions

Carbon dioxide (CO<sub>2</sub>) emissions from fossil fuels and industry<sup>1</sup>. Land-use change is not included.

Our World  
in Data




Data source: Global Carbon Budget (2023); Population based on various sources (2023)  
OurWorldinData.org/co2-and-greenhouse-gas-emissions | CC BY

1. **Fossil emissions:** Fossil emissions measure the quantity of carbon dioxide (CO<sub>2</sub>) emitted from the burning of fossil fuels, and directly from industrial processes such as cement and steel production. Fossil CO<sub>2</sub> includes emissions from coal, oil, gas, flaring, cement, steel, and other industrial processes. Fossil emissions do not include land use change, deforestation, soils, or vegetation.


# The basics: Green house effect

## Black body radiation of the sun and Earth

- Sun at 6000°C, Earth at 15°C → Sun radiates in the visible, Earth radiates in the infra-red




Credit: By 4C - Own work based on JPG version Curva Planck TT.jpg, CC BY-SA 3.0, <https://commons.wikimedia.org/w/index.php?curid=1017820>



→ If atmosphere did not re-absorb Earth's emissions, surface temperature on Earth around -18°C!

# The basics: Green house effect


## Re-absorption done by greenhouse gases



### Main greenhouse gases

→ Historic concentrations in last ~800k years

- Water vapour (strongly varying)
- Carbon dioxide (CO<sub>2</sub>) → ~240ppm
- Methane (CH<sub>4</sub>) → ~500ppb
- Nitrous oxide (N<sub>2</sub>O) → ~250ppb
- Fluorinated gases: Hydrofluorocarbons, perfluorocarbons, sulfur hexafluoride, nitrogen trifluoride → Historically not existing




<https://www.epa.gov/ghgemissions/overview-greenhouse-gases>

Historic contributions (pre-industrial age) read off from figures from: <https://www.epa.gov/climate-indicators/climate-change-indicators-atmospheric-concentrations-greenhouse-gases>

# Improving the predictions of Earth's surface temperature

Nobel prize 2021 - *"for groundbreaking contributions to our understanding of complex physical systems"*


- 1967: Syukuro Manabe: Adding convection and latent heat



→ If the cause for warming was increased solar radiation, the entire atmosphere should have warmed up

→ Hence, cause is increased CO<sub>2</sub> levels

With higher CO<sub>2</sub> levels ...



<https://www.nobelprize.org/prizes/physics/2021/popular-information/>




# Improving the predictions of Earth's surface temperature

Nobel prize 2021 - *"for groundbreaking contributions to our understanding of complex physical systems"*

- Around 1980: Klaus Hasselmann: Building a stochastic climate model


- How to make reliable climate predictions, while weather forecasts are notoriously imprecise in the long-term?
- Treatment of weather as rapidly changing noise
- Human impact separated out by properties of noise and signals → Unique fingerprints
  - Solar radiation
  - Volcanic particles
  - Levels of greenhouse gases
  - Human impact



<https://www.nobelprize.org/prizes/physics/2021/popular-information/>

# Climate simulation


In 2007



<https://data.giss.nasa.gov/modelE/sc07/>

# Impacts attributed to human influence

## Driven by changes in multiple physical climate conditions




[https://www.ipcc.ch/report/ar6/syr/downloads/report/IPCC\\_AR6\\_SYR\\_SPM.pdf](https://www.ipcc.ch/report/ar6/syr/downloads/report/IPCC_AR6_SYR_SPM.pdf)  
[https://report.ipcc.ch/ar6syr/pdf/IPCC\\_AR6\\_SYR\\_LongerReport.pdf](https://report.ipcc.ch/ar6syr/pdf/IPCC_AR6_SYR_LongerReport.pdf)

# Generations affected by climate change

## Considering the different scenarios

c) The extent to which current and future generations will experience a hotter and different world depends on choices now and in the near-term




[https://www.ipcc.ch/report/ar6/syr/downloads/report/IPCC\\_AR6\\_SYR\\_SPM.pdf](https://www.ipcc.ch/report/ar6/syr/downloads/report/IPCC_AR6_SYR_SPM.pdf)



# Risk of species losses


**Risk of species losses**  
Percentage of animal species and seagrasses exposed to potentially dangerous temperature conditions<sup>1, 2</sup>



# Study of remaining carbon budget newer than IPCC report


Lamboll et. Al., Nature Climate Change 2023

## For 1.5°C increase max



Reduced complexity climate model for non-CO<sub>2</sub> emissions

## For 2.0°C increase max




Additional simple climate model calibrated for use in the latest IPCC report

→ For 50% chance at 1.5°C (2.0°C) global warming, only ~250GtCO<sub>2</sub> (~1220GtCO<sub>2</sub>) remaining budget, as of January 2023!  
→ With 40GtCO<sub>2</sub> emissions in 2022: net zero CO<sub>2</sub> emissions needed by 2029 (2054)!


# IPCC report: Mitigation potentials

## Cost estimates of different mitigation options



- Energy and food production with large impacts
- Take a closer look at these two next

[https://report.ipcc.ch/ar6syrr/pdf/IPCC\\_AR6\\_SYR\\_LongerReport.pdf](https://report.ipcc.ch/ar6syrr/pdf/IPCC_AR6_SYR_LongerReport.pdf)




Net lifetime cost of options:


- Costs are lower than the reference
- 0–20 (USD per tCO<sub>2</sub>-eq)
- 20–50 (USD per tCO<sub>2</sub>-eq)
- 50–100 (USD per tCO<sub>2</sub>-eq)
- 100–200 (USD per tCO<sub>2</sub>-eq)
- Cost not allocated due to high variability or lack of data

# Greenhouse gases

## CO<sub>2</sub>




## Methane




<https://www.epa.gov/climate-indicators/climate-change-indicators-atmospheric-concentrations-greenhouse-gases>

# Greenhouse gases

## Nitrous Oxide



## Halogenated Gases



<https://www.epa.gov/climate-indicators/climate-change-indicators-atmospheric-concentrations-greenhouse-gases>



# Gas emissions as main driver of CO<sub>2</sub> footprint

## Global warming potential (GWP) of gas


- How much energy will be absorbed by 1t of the gas in 100 (500) years compared to 1t of CO<sub>2</sub>?

## Gases used at CERN

- Have significant GWPs > 1000 or even 10000

GROUP	GASES	tCO <sub>2</sub> e 2019	tCO <sub>2</sub> e 2020
PFC	CF <sub>4</sub> , C <sub>2</sub> F <sub>6</sub> , C <sub>3</sub> F <sub>8</sub> , C <sub>4</sub> F <sub>10</sub> , C <sub>6</sub> F <sub>14</sub>	43277	45678
HFC	CHF <sub>3</sub> (HFC-23), C <sub>2</sub> H <sub>2</sub> F <sub>4</sub> (HFC-134a), HFC-404a, HFC-407c, HFC-410a, HFC R-422D, HFC-507	17540	34899
Other F-gases	SF <sub>6</sub> , NOVEC, R1234ze	3840	5377
CO <sub>2</sub>	CO <sub>2</sub>	13512	13046
<b>TOTAL SCOPE 1</b>		<b>78169</b>	<b>98997</b>

GWP <sub>100</sub> (*)	GWP <sub>500</sub> (*)
7390, 12200, 8830, 8860, 9300	11200, 18200, 12500, 12500, 13300
14800, 1430	12000, 435
22800	32600
1	



Note: C<sub>4</sub>H<sub>10</sub> = Butane: GWP(100years) = 4.0 (\*)

→ Already very small leaks have a major impact

→ Circled gases are also used at CERN

(\*) <https://www.ipcc.ch/site/assets/uploads/2018/02/ar4-wg1-chapter2-1.pdf>


<https://www.epa.gov/climate-indicators/climate-change-indicators-atmospheric-concentrations-greenhouse-gases>

# Hands-on: Fixing leaks in ATLAS

ATL-MUON-  
PROC-2023-  
001

## Gas leaks in the ATLAS muon system

- Plastic connectors of the gas flow lines to the Resistive Plate Chambers (RPCs) → Tend to develop leaks
- 8000 potentially leaky connection points in ATLAS RPCs → Often difficult to reach → Break faster than can be repaired
- Gas mixture in RPCs:  $\text{C}_2\text{H}_2\text{F}_4 + \text{iso-C}_4\text{H}_{10} + \text{SF}_6$  → GWP ~ 1400 → Studies with replacing gas mixture not trivial!
  - 1l of RPC mixture ~ 5-6kg  $\text{CO}_2$ -eq. (\*) → Loss of ~1000l/h  
→ If constant throughout the year: ~44k-53k  $\text{tCO}_2$ -eq./year emissions  
→ ~20-23% of 2018 emissions by CERN (own estimate)
- Campaign in ATLAS with new repair technique and teams of volunteers to fix leaks during end-of-year shutdowns
  - First test campaign early 2023: Reduction of RPC losses by 23%!
  - Needs follow-up in further shutdowns




(\*) Based on main component:  $\text{C}_2\text{F}_2\text{F}_4$  → [Conversion of l to kg](#) → Convert to  $\text{CO}_2$ -eq. by multiplying with [GWP](#) for HFC-134(a)

# Relation to SDGs

(b) Climate responses and adaptation options have benefits for ecosystems, ethnic groups, gender equity, low-income groups and the Sustainable Development Goals

Relations of sectors and groups at risk (as observed) and the SDGs (relevant in the near-term, at global scale and up to 1.5°C of global warming) with climate responses and adaptation options



Footnotes: <sup>1</sup> The term response is used here instead of adaptation because some responses, such as retreat, may or may not be considered to be adaptation. <sup>2</sup> Including sustainable forest management, forest conservation and restoration, reforestation and afforestation. <sup>3</sup> Migration, when voluntary, safe and orderly, allows reduction of risks to climatic and non-climatic stressors. <sup>4</sup> The Sustainable Development Goals (SDGs) are integrated and indivisible, and efforts to achieve any goal in isolation may trigger synergies or trade-offs with other SDGs. <sup>5</sup> Relevant in the near-term, at global scale and up to 1.5°C of global warming.

# What barriers exist for getting involved?

## Psychological barriers to climate action



Example: Moral barrier


→ Broad categories of morality



Avoiding harm



Fairness



Loyalty



Respecting authority



Purity

→ Most often addressed by climate crisis communication

→ Leaves out a huge part of the population

Left-leaning

Political spectrum


Right-leaning

→ Need to adjust messaging to include entire population!

Source: [Presentation by Prof. Brosch, at CERN and the environment workshop, Oct 2022](#)

# HL-LHC operation schedule

## Start of HL-LHC with Run 4



Last update: September 24



# Details on travel

---

## Long-distance buses vs. Long-distance trains

Source of Emission	Emission Factor	
Long-distance Buses	0.031	kgCO <sub>2</sub> e/km
Long-distance Trains	0.031	kgCO <sub>2</sub> e/km
Personal Car	0.17	kgCO <sub>2</sub> e/km
Flights within Europe	130	kgCO <sub>2</sub> e/h
Transcontinental Flights	170	kgCO <sub>2</sub> e/h
Hotel room	12	kgCO <sub>2</sub> e/night
Event venue	0.19	kgCO <sub>2</sub> e/day

→ Why the same emission factor?

- By chance! → For UBA numbers from 2022 (<https://www.umweltbundesamt.de/themen/verkehr/emissionsdaten>)
  - Tank-to-wheel (TTW) for buses much higher than for trains, i.e. running a bus has higher emissions than a train
  - Compensated by well-to-tank (WTT) for trains and infrastructure, i.e. the extraction of the fuel (using German conventional electricity mix) + the building of the infrastructure (rails, etc.) more costly for trains than for buses